

The Atomic Limit of the Boson Fermion Model

T. Domanski[†], J. Ranninger and J. M. Robin

Centre de Recherches sur les Très Basses Températures, Laboratoire Associé à l'Université

Joseph Fourier, Centre National de la Recherche Scientifique,

BP 166, 38042, Grenoble Cédex 9, France

(February 1, 2008)

Abstract

The Boson-Fermion model, describing a mixture of hybridized localized Bosons and itinerant Fermions on a lattice, is known to exhibit spectral properties for the Fermions which upon lowering the temperature develop into a three pole structure in the vicinity of the Fermi level. These spectral features go hand in hand with the opening of a pseudogap in the density of states upon approaching the critical temperature T_c when superconductivity sets in. In the present work we study this model, in the atomic limit where the three pole structure arises naturally from the local bonding, anti-bonding and non-bonding states between the Bosons and Fermions.

PACS numbers: 79.60.-i, 74.25.-q, 74.72.-h

The Boson Fermion model presents one of the possible scenarios which are presently widely discussed in connection with the phenomenon of High T_c superconductivity [1]. This model treats localized Bosons (such as bi-polarons) which are locally hybridized with pairs of itinerant Fermions (band electrons). On the basis of this model one envisages the opening of a pseudogap in the density of states of the electrons as the temperature descends below a certain characteristic temperature T^* [2] and approaches T_c from above. The opening of the pseudogap has been attributed to a precursor effect of superfluid phase fluctuations [3] with the amplitude of the order parameter remaining finite. Near the Fermi level this is tantamount to the existence of resonance two-particle states into which most of the electrons are engaged. The physical implications of these one-particle features on the thermodynamic, transport and magnetic properties demonstrate this behaviour, which radically differs from that expected for classical Fermi liquids [4]. The deviations from Fermi liquid properties are manifest in a complex three-pole structure of the one-particle electron Green's function [3]. These findings have been obtained by solving selfconsistently the equations for the Boson and Fermion selfenergies using a diagrammatic conserving approximation, a procedure from which the underlying physics could be inferred.

In this paper, we show that some of these physical interpretations can indeed be put on a firm basis when studying the atomic limit of this model. In this case, the local interaction between a hard core Boson and a pair of Fermion (having opposite spins) on each site of the lattice can be diagonalized exactly. The kinetic part of the Fermions is subsequently treated in an approximation à la Hubbard.

The Boson-Fermion hamiltonian in the Wannier representation is given by [5]

$$\begin{aligned}
H = & (zt - \mu) \sum_{j,\sigma} c_{j,\sigma}^\dagger c_{j,\sigma} - t \sum_{j,\delta,\sigma} c_{j+\delta,\sigma}^\dagger c_{j,\sigma} + (\Delta_B - 2\mu) \sum_j b_j^\dagger b_j \\
& + v \sum_j \left[c_{j,\uparrow}^\dagger c_{j,\downarrow}^\dagger b_j + b_j^\dagger c_{j,\downarrow} c_{j,\uparrow} \right], \tag{1}
\end{aligned}$$

where the sum over δ runs over the z nearest neighbours. μ denotes the chemical potential, Δ_B the localized Boson level, t the hopping integral for the Fermions and v the strength

of the local hybridization. The Fermionic operators $c_{j,\sigma}^{(\dagger)}$ anticommute while the bosonic operators $b_j^{(\dagger)}$ commute for different sites and anticommute for the same site. The Hilbert space for each site is then the direct product of the Fermionic and Bosonic states:

$$\{ |0\rangle, |\uparrow\rangle, |\downarrow\rangle, |\uparrow\downarrow\rangle \} \otimes \{ |0\rangle, |1\rangle \}. \quad (2)$$

The local interaction term mixes the states $|\uparrow\downarrow\rangle \otimes |0\rangle$ and $|0\rangle \otimes |1\rangle$. The corresponding bonding and anti-bonding eigenstates are

$$\begin{cases} |B^-\rangle = U |\uparrow\downarrow\rangle \otimes |0\rangle - V |0\rangle \otimes |1\rangle \\ |B^+\rangle = V |\uparrow\downarrow\rangle \otimes |0\rangle + U |0\rangle \otimes |1\rangle \end{cases} \quad (3)$$

with the associated eigenvalues $\varepsilon_{\pm} = (2\varepsilon_0 + E_0)/2 \pm \gamma$.

$$U^2 = \frac{1}{2} \left(1 - \frac{2\varepsilon_0 - E_0}{2\gamma} \right), \quad V^2 = \frac{1}{2} \left(1 + \frac{2\varepsilon_0 - E_0}{2\gamma} \right), \quad UV = \frac{v}{2\gamma} \quad (4)$$

and $\varepsilon_0 = zt - \mu$, $E_0 = \Delta_B - 2\mu$ and $\gamma = \sqrt{(\varepsilon_0 - E_0/2)^2 + v^2}$. In the atomic limit the one-particle Fermionic local Green's function $G_{Loc}(\tau) = -\langle T [c_{j,\uparrow}(\tau) c_{j,\uparrow}^\dagger] \rangle$ is then given by

$$G_{Loc}(i\omega_n) = \frac{Z_f}{i\omega_n - \varepsilon_0} + \frac{1 - Z_f}{i\omega_n - v^2 / (i\omega_n + \varepsilon_0 - E_0)} \quad (5)$$

where Z_f is the temperature dependent factor

$$Z_f = \frac{1 + e^{-\beta\varepsilon_0} + e^{-\beta(\varepsilon_0 + E_0)} + e^{-\beta(2\varepsilon_0 + E_0)}}{1 + 2e^{-\beta\varepsilon_0} + 2e^{-\beta(\varepsilon_0 + E_0)} + e^{-\beta(2\varepsilon_0 + E_0)} + e^{-\beta(\varepsilon_0 + E_0/2 - \gamma)} + e^{-\beta(\varepsilon_0 + E_0/2 + \gamma)}} \quad (6)$$

and $\beta = 1/k_B T$ denotes the inverse of the temperature. This Green's function contains three poles with spectral weights Z_f , $U^2(1 - Z_f)$ and $V^2(1 - Z_f)$. The first term in the expression for this Green's function, Equ.(5), determines the contribution from the non-bonding one-particle Fermionic states, while the second term is reminiscent of Bogoliubov type excitations and contains the effect of hybridized two-particle bonding and respectively anti-bonding states, $|B^-\rangle$ and $|B^+\rangle$. In the remainder of this paper we consider the

symmetric case i.e., $\varepsilon_0 = 0$ and $E_0 = 0$ for which $\langle c_{j,\sigma}^\dagger c_{j,\sigma} \rangle = \langle b_j^\dagger b_j \rangle = 1/2$. We then obtain $Z_f = 2/(3 + \cosh(\beta v))$ and are thus left with only one parameter, namely βv . The non-bonding component, corresponding to the single particle pole ε_0 has a finite spectral weight for any $T \neq 0$, while for $T = 0$ it vanishes.

Let us now generalize the above atomic Green's function to the case of a lattice. The corresponding Green's function $G(\mathbf{k}, i\omega_n)$ is obtained by adding à la Hubbard the kinetic term which is diagonal in $\mathbf{k}$ -space:

$$G(\mathbf{k}, i\omega_n) = \frac{1}{[G_{Loc}(i\omega_n)]^{-1} - t_{\mathbf{k}}} \quad (7)$$

where $t_{\mathbf{k}}$ is the Fourier's transformation of the hopping term [6]. The three-pole structure is then repercutated from the atomic Green's function onto the lattice Green's function whose dispersion branches are illustrated in Fig.(1) and which resemble those previously derived by us on the basis of a selfconsistent diagrammatic approach [3]. Using a free parabolic density of states of the form

$$\rho_0(\varepsilon) = \frac{8}{\pi D} \sqrt{(\varepsilon/D + 1/2)(\varepsilon/D - 1/2)} \quad , \quad -D/2 \leq \varepsilon \leq D/2 \quad (8)$$

where D denoted the bandwidth of the Fermions $D = 2zt$, we obtain a one-particle Green's function, which has the following two-pole structure:

$$G(\mathbf{k}, i\omega_n) = \frac{1}{2} \left(1 + \frac{t_{\mathbf{k}}}{4\gamma_{\mathbf{k}}} \right) \frac{1}{i\omega_n - t_{\mathbf{k}}/2 - \gamma_{\mathbf{k}}} + \frac{1}{2} \left(1 - \frac{t_{\mathbf{k}}}{4\gamma_{\mathbf{k}}} \right) \frac{1}{i\omega_n - t_{\mathbf{k}}/2 + \gamma_{\mathbf{k}}} \quad (9)$$

provided the temperature is exactly equal to zero. With $\gamma_{\mathbf{k}} = \sqrt{t_{\mathbf{k}}^2/4 + v^2}$, this form is reminiscent of the one-particle Green's function exhibiting a Bogoliubov like spectrum as in the case of a BCS superconductor. The corresponding density of states

$$\rho(\varepsilon) = \rho_0(\varepsilon - v^2/\varepsilon) \left[\theta(\varepsilon - E_1^-) \theta(E_1^+ - \varepsilon) + \theta(\varepsilon - E_2^-) \theta(E_2^+ - \varepsilon) \right] \quad (10)$$

with $E_1^\pm = \pm \frac{D}{4} + \frac{1}{2} \sqrt{\left(\frac{D}{2}\right)^2 + 4v^2}$ and $E_2^\pm = \pm \frac{D}{4} - \frac{1}{2} \sqrt{\left(\frac{D}{2}\right)^2 + 4v^2}$ thus shows a gap of size $E_G = 2E_1^-$ at the Fermi level $\varepsilon = 0$ for $T = 0$. Our previous study of the Boson Fermion model clearly showed indications for such a gap in form of a dip in the density of states which deepened upon lowering the temperature [3].

For any finite temperature $T \neq 0$ the density of states $\rho(\varepsilon)$ at the Fermi level turns out to be finite and moreover we find $\rho(0) = \rho_0(0)$. What changes with temperature is the spectral weight Z_f of the normal contribution of the single particle excitations. It decreases to zero as the temperature decreases which is manifest in a gradual narrowing of the peak in $\rho(\varepsilon)$ corresponding to the non-bonding one-particle state, centered around the Fermi level as illustrated in Fig.(2). This behaviour can be derived from the expression for the one-particle Green's function which in the limit $|\varepsilon| \ll v$ can be derived from the correspondingly approximated spectral function

$$A(\mathbf{k}, \varepsilon) = -\frac{1}{\pi} \text{Im} G(\mathbf{k}, \varepsilon + i0) \simeq -\frac{1}{\pi} \text{Im} \frac{Z_f}{\varepsilon + i0 - Z_f t_{\mathbf{k}}} \quad (11)$$

the density of states is then obtained as

$$\rho(\varepsilon) = \sum_{\mathbf{k}} A(\mathbf{k}, \varepsilon) = Z_f \int_{-D/2}^{+D/2} d\omega \rho_0(\omega) \delta(\varepsilon - Z_f \omega) = \rho_0(\varepsilon/Z_f) \quad (12)$$

from which we can see the narrowing down of the peak associated in the non-bonding component of the spectrum which scales the free density of states ρ_0 with the temperature dependent factor Z_f .

From the present discussion we see that the opening of the pseudogap obtained in our previous selfconsistent treatment of the Boson-Fermion model [3] is clearly associated with the disappearance of the non-bonding component of the spectrum. As the temperature is decreased its spectral weight is transferred into the contributions coming from the hybridized Boson-Fermion pairs in bonding and anti-bonding states. These contributions to the density of states are centered around $\varepsilon_{\pm}$ (for which moreover we have $\max \rho(\varepsilon) = \rho_0(0)$ for $\varepsilon = \varepsilon_{\pm}$, as can be seen directly from Eq.(10)). The temperature variation of the partial density of states, coming from those non-bonding contributions, is illustrated in Fig.(3). We notice that it scales with Z_f and that it shows a marked saturation above a certain characteristic temperature which we identify with the temperature T^* where a pseudogap opens up in the density of states and which was previously obtained by us using a fully selfconsistent treatment of the Boson-Fermion model [3,4]. The saturation above T^* , which is of the order

of the coupling constant v , corresponds to a blocking of the Boson - Fermion-pair exchange mechanism.

The present treatment of the Boson-Fermion model captures the underlying physics of the opening of a pseudogap in the density of states due to strong local two-particle bonding and anti-bonding resonances which increase in spectral weight as the temperature decreases. This can be seen from the superfluid pair susceptibility for the Bosons $\chi_B = -B(i\omega_n = 0)$ and for the Fermions $\chi_F = -G_{FF}(i\omega_n = 0)$, which tends to a finite value as T decreases to zero. Given the onsite Green's functions $B(\tau) = -\langle T[b_j(\tau)b_j^\dagger] \rangle$ and $G_{FF}(\tau) = -\langle T[c_{j,\downarrow}(\tau)c_{j,\uparrow}(\tau)c_{j,\uparrow}^\dagger c_{j,\downarrow}^\dagger] \rangle$, the superfluid pair susceptibilities for the Bosons and Fermions are determined by

$$\chi_B = \frac{1}{v} \frac{\beta v + \sinh(\beta v)}{3 + \cosh(\beta v)}, \quad \chi_F = \frac{1}{v} \frac{\sinh(\beta v)}{3 + \cosh(\beta v)}. \quad (13)$$

In the limit $\beta v \Rightarrow \infty$ we hence obtain $\chi_B \Rightarrow \frac{1}{v}$ and $\chi_F \Rightarrow \frac{1}{v}$, instead of $\frac{1}{T}$ as for the uncoupled system, which indicates the strong two-particle resonance behaviour without any superfluid coherence. A further indication for strong pairing in the normal state comes from the expectation value of the number of pairs

$$n_p = \langle c_{j,\uparrow}^\dagger c_{j,\downarrow}^\dagger c_{j,\downarrow} c_{j,\uparrow} \rangle = \frac{1 + \cosh(\beta v)}{6 + 2\cosh(\beta v)} \quad (14)$$

In the limit of the temperature going to zero, i.e., $\beta v \Rightarrow \infty$ we obtain $n_p \Rightarrow \frac{1}{2}$, thus showing a significant enhancement over its free particle value, which is $\frac{1}{4}$.

These results, obtained in the atomic limit, clearly demonstrate that the pseudogap in the DOS of the Fermions is linked to the strong local pair correlations in the normal state with any global phase coherence being absent. A more refined treatment of those local resonant states (such as given in our previous study [3] which includes spatial correlations of the two-particle resonant states) leads in the normal state to a "PAIR LIQUID" with fluctuations of a condensate of those two-particle states, which ultimately changes the nature of the gap leading to a superconducting ground state.

I. ACKNOWLEDGEMENT

T.D. would like to acknowledge a grant from the French ministry of High Education and Scientific Research and support from the Polish Committee of Scientific Research under the project No. 2P03B03111.

† On leave of absence from the Institute of Physics, Maria Curie Skłodowska University, Lublin, Poland.

REFERENCES

- [1] I.O. Kulik, Int. J. Mod. Phys. B **1**, 851 (1988); G.M. Eliashberg, JETP Lett. **46**, 581 (1987); L. Ioffe, A.I. Larkin, Yu.N. Ovchinnikov and L. Yu, Int. J. Mod. Phys. B **3**, 2065 (1989); J.R. Friedberg and T.D. Lee, Phys. Rev. B **40**, 6745 (1989); Y.Bar Yam, Phys. Rev. B **43**, 359 (1991); B.R. Alascio and C.R. Proetto, Solid State Commun. **75**, 217 (1990); H.Eschrig and S.L. Drechsler, Physica C **173**,80 (1991), D.M. Eagles, Solid State Commun. **80**,823 (1991).
- [2] B. Batlogg, Physica C **235 - 240**,130 (1991).
- [3] J. Ranninger, J.M. Robin and M. Eschrig, Phys. Rev. Lett. **74**, 4027 (1995); J. Ranninger and J. M. Robin, Solid State Commun. **98**, 559 (1996).
- [4] J. Ranninger and J. M. Robin, Phys. Rev. **B53**, R11961 (1996).
- [5] For a physical justification of this model see for instance J. Ranninger and S.Robaszkiewicz, Physica B **135**, 468 (1985) where this model was proposed for the first time and J. Ranninger and J. M. Robin, Physica **C253**, 279 (1995).
- [6] J. Hubbard, Proc. Roy. Soc. **A276**, 238 (1963); S. Doniach and E. H. Sondheimer, *Green's functions for Solid State Physicists*, chap. 8, BENJAMIN CUMMINGS (1974).

FIGURES

Fig.1 The dispersion of the three poles of the one-particle Green's function in the atomic limit for $v = 0.2$.

Fig.2 The density of states $\rho(\varepsilon)$ for four different temperatures $T = 0.1$ Fig.(2a), $T = 0.05$ Fig.(2b), $T = 0.001$ Fig.(2c) and $T = 0$ Fig.(2d) which clearly shows the disappearance of the non-bonding component as the temperature is decreased.

Fig.3 Temperature variation of the partial integrated density $I(T)$ of states arising from the contribution of the non-bonding states.











